

# ISOLATION AND IDENTIFICATION OF FUNGI FROM RHIZOSPHERE IN THE VICINITY OF THE INSTITUTE OF AGRICULTURAL SCIENCES UNIVERSITY OF THE PUNJAB, LAHORE PAKISTAN

ATIF M, MANZOOR MT\*, SARWAR M, RAFAQAT N

Institute of Agricultural Sciences, University of the Punjab, Lahore, Punjab, Pakistan

\*Correspondence author email address: [tariq.iags@pu.edu.pk](mailto:tariq.iags@pu.edu.pk)

(Received, 4<sup>th</sup> January 2019, Revised 23<sup>rd</sup> December 2019, Published 25<sup>th</sup> December 2019)

**Abstract:** Rhizosphere and endorhizosphere microbial colonization aid in phytoremediation, carbon sequestration, and plant development. According to several studies, the microbiomes of various plant species' genotypes and even plant species might vary somewhat. Both pathogenic and symbiotic fungi are associated with the rhizosphere, but the specific fungal communities depend on various conditions, including plant growth and soil type. The present study was conducted to isolate and identify fungus species from vicinity of the Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan and identified *Aspergillus*, *Mucor*, *Alternaria*, *Penicillium*, four fungal species from all of the 10 samples.

[Citation: Atif, M., Manzoor, M.T., Sarwar, M., Rafiqat, N. (2019). Isolation and identification of fungi from rhizosphere in the vicinity of the Institute of Agricultural Sciences University of the Punjab, Lahore Pakistan. *Bull. Biol. All. Sci. Res.* 4: 30. doi: <https://doi.org/10.54112/bbasr.v2019i1.30>]

**Keywords:** Rhizosphere, phytoremediation, pathogenic, symbiotic, fungi

## Introduction

The rhizosphere is an extremely favorable environment for various microorganisms' growth, activity, and metabolism. The rhizosphere microflora may be extensively counted through microscopic, cultural, and biochemical methods. For the earth to remain healthy and to support life there, a greater knowledge of the rhizosphere is essential. A small but focused effort is now under work to boost the production potential of major food crops and satisfy the predicted increase in global food demand over the next 50 years (Zhang *et al.*, 2010). The cortex and endodermis's "free space" between cells can be occupied by cations and bacteria in the endorhizosphere (apoplastic space). The medial zone immediately around the root, which includes the root epidermis and mucilage, is known as the rhizoplane. The ectorhizosphere extends from the rhizoplane into the bulk soil and is the outermost zone. The rhizosphere is not an area with a defined size or shape, as expected, given the intrinsic complexity and diversity of plant root systems. Instead, it comprises a gradient of chemical, biological, and physical qualities that vary radially and longitudinally down the root (Hartmann *et al.*, 2008). A generic name used to describe the symbiotic association between a soil fungus and plant root is mycorrhizae, which derives from the Greek words for fungus and root. Mycorrhizal partnerships, which occur in around 80% of angiosperms and all gymnosperms, are widespread

and largely nonselective, in contrast to rhizobia and their legume partners (Wilcox, 1991). Mycorrhiza aid plants in drawing water, phosphorus, and other micronutrients from the soil (such as Zn and Cu) in exchange for receiving food (carbon) from the plant (El-Morsy, 1999). Rhizosphere fungi naturally create a beneficial symbiotic connection with most plants, giving their hosts nitrogen and phosphorus in exchange. This interaction can considerably increase the efficiency with which plants absorb soil nutrients (Rilling *et al.*, 2018).




## Material and Methods

10 trees were selected present in the vicinity of the Institute of Agricultural sciences, University of the Punjab, Lahore, Pakistan including.

1. *Syzygium cumini* (Jaman)
2. *Mangifera indica* (Mango)
3. *Phoenix dactylifera* (Date palm)
4. *Albizia lebbek* (shareen)
5. *Ziziphus mauritiana* (Bair)
6. *Cassia fistula* (Amaltas)
7. *Acacia nilotica* (kiker)
8. *Dalburgia sisso* (shesham)
9. *Bauhinia variegata* (kachnar)
10. *Capparis decidua* (karir)

The rhizosphere soil was collected in sampling bags, and each soil bag was labelled with the tree name and date of collection. Small rootlets were also collected, adhering to rhizospheric soil. Malt Agar

media is used for growth to isolate and purify fungi. Soil collected from rhizosphere was mixed with water and diluted with water using the serial dilution method. Inoculum was taken by the dropper from the last test tube, put in the centre of the Malt Ager plate, and then spread by the spreader on the whole plate, and the lid was closed. This was repeated for all ten samples. All Media plates were incubated at 25-27°C for 5-7 days.



#### Isolation of fungal microbes from Ten trees

After isolation, microscopy was used to identify fungi in rhizosphere of 10 trees. The Inoculum needle was heated for sterilization purposes, and the inoculum was taken and placed in the center of the slide and stain/ simple water was poured with the dropper. Slides were observed under the microscope. And

identified with their characteristic fruiting bodies and hyphae. *Aspergillus*, *Mucor*, *Alternaria*, *Penicillium*, these four fungal species, were identified in all 10 samples.






Fig D, *Mucor*, Fig E, *Penicillium*, Fig F, *Alternaria*, Fig G, *Aspergillus*

#### Conflict of interest

The authors declared the absence of a conflict of interest.

#### References

El-Morsy, E. M. (1999). Microfungi from the ectorrhizosphere-rhizoplane zone of different halophytic plants from the Red Sea coast of

- Egypt. *Mycologia*, **91**, 228-236 DOI <https://doi.org/10.2307/3761367>
- Hartmann A., Rothballer M., and M. Schmid. Lorenz Hiltner. (2008). A pioneer in rhizosphere microbial ecology and soil bacteriology research. *Plant Soil*, **312**, 7-14. DOI <https://doi.org/10.1007/s11104-007-9514-z>
- Rilling, J., Acuña, J., Nannipieri, P., Cassan, F., Maruyama, F., Jorquera, M. (2018). Current opinion and perspectives on the methods for tracking and monitoring plant growth-promoting bacteria. *Soil Biology and Biochemistry*, **130**, 205-219. DOI <https://doi.org/10.1016/j.soilbio.2018.12.012>
- Wilcox, H.E. (1991). *Mycorrhizae*. In: Waisel Y, Eshel A, Kafkaki U, eds. Plant roots: the hidden half. New York, USA: Marcel Dekker.
- Zhang, F., Shen, J., Zhang, J., Zuo, Y. Li, L., Chen, X. (2010). Rhizosphere processes and management for improving nutrient use efficiency and crop productivity: implications for China. *Advances in Agronomy*. **107**, 1-32. DOI [https://doi.org/10.1016/S0065-2113\(10\)07001-X](https://doi.org/10.1016/S0065-2113(10)07001-X)



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution, and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third-party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. © The Author(s) 2019